

# Murine T-box transcription factor *Tbx20* acts as a repressor during heart development, and is essential for adult heart integrity, function and adaptation

Fiona A. Stennard<sup>1</sup>, Mauro W. Costa<sup>1,2</sup>, Donna Lai<sup>1</sup>, Christine Biben<sup>1</sup>, Milena B. Furtado<sup>1</sup>, Mark J. Solloway<sup>1</sup>, David J. McCulley<sup>3</sup>, Christiana Leimena<sup>1</sup>, Jost I. Preis<sup>1</sup>, Sally L. Dunwoodie<sup>1,4</sup>, David E. Elliott<sup>1,\*</sup>, Owen W. J. Prall<sup>1</sup>, Brian L. Black<sup>3</sup>, Diane Fatkin<sup>1,4</sup> and Richard P. Harvey<sup>1,4,†</sup>

<sup>1</sup>Victor Chang Cardiac Research Institute, St Vincent's Hospital, 384 Victoria Street, Darlinghurst 2010, New South Wales, Australia

<sup>2</sup>Laboratório de Cardiologia Celular e Molecular, Instituto de Biofísica Carlos Chagas Filho, Universidade Federal do Rio de Janeiro, 20941-000, Brazil

<sup>3</sup>Cardiovascular Research Institute, University of California, San Francisco, CA 94143-0130, USA

<sup>4</sup>Faculties of Medicine and Life Sciences, University of New South Wales, Kensington 2056, New South Wales, Australia

\*Present address: The Wellcome Trust/Cancer Research UK, Gurdon Institute of Cancer and Developmental Biology, University of Cambridge, Tennis Court Road, Cambridge CB2 1QR, UK

†Author for correspondence (e-mail: r.harvey@victorchang.unsw.edu.au)

Accepted 21 February 2005

Development 132, 2451-2462  
Published by The Company of Biologists 2005  
doi:10.1242/dev.01799

## Summary

The genetic hierarchies guiding lineage specification and morphogenesis of the mammalian embryonic heart are poorly understood. We now show by gene targeting that murine T-box transcription factor *Tbx20* plays a central role in these pathways, and has important activities in both cardiac development and adult function. Loss of *Tbx20* results in death of embryos at mid-gestation with grossly abnormal heart morphogenesis. Underlying these disturbances was a severely compromised cardiac transcriptional program, defects in the molecular pre-pattern, reduced expansion of cardiac progenitors and a block to chamber differentiation. Notably, *Tbx20*-null embryos showed ectopic activation of *Tbx2* across the whole heart myogenic field. *Tbx2* encodes a transcriptional repressor normally expressed in non-chamber myocardium, and in the atrioventricular canal it has been proposed to inhibit chamber-specific gene expression

through competition with positive factor *Tbx5*. Our data demonstrate a repressive activity for *Tbx20* and place it upstream of *Tbx2* in the cardiac genetic program. Thus, hierarchical, repressive interactions between *Tbx20* and other T-box genes and factors underlie the primary lineage split into chamber and non-chamber myocardium in the forming heart, an early event upon which all subsequent morphogenesis depends. Additional roles for *Tbx20* in adult heart integrity and contractile function were revealed by in-vivo cardiac functional analysis of *Tbx20* heterozygous mutant mice. These data suggest that mutations in human cardiac transcription factor genes, possibly including *TBX20*, underlie both congenital heart disease and adult cardiomyopathies.

Key words: T-box, *Tbx20*, Heart, *Nkx2-5*, *Tbx2*, Chamber myocardium, Dilated cardiomyopathy, mice

## Introduction

Cardiac development in mammals is guided by an ancient and conserved genetic program (Cripps and Olson, 2002; Harvey, 2002; Zaffran and Frasch, 2002). However, how the cardiac program unfolds, the specific relationships between patterning events and the transcription factor hierarchy, and how cardiomyocyte function impacts on heart form remain poorly understood.

Of 18 T-box factor genes identified in mammals, at least six of them (*Tbx1/2/3/5/18/20*) are expressed in the developing heart (Plageman and Yutzey, 2004). T-box proteins are characterized by the presence of a sequence-specific DNA-binding domain called the T-box (Smith, 1999). During embryogenesis, T-box genes are expressed in restricted and sometimes overlapping domains throughout gastrulation and/or organogenesis, and in some cases roles in controlling

cell fate and migration have been demonstrated (Chapman and Papaioannou, 1998; Naiche and Papaioannou, 2003; Russ et al., 2000). T-box factors can act up- or downstream of signaling factors of the TGF- $\beta$  (Suzuki et al., 2004), fibroblast growth factor (Brown et al., 2004; Hu et al., 2004; Sakiyama et al., 2003; Yamagishi et al., 2003), sonic hedgehog (Suzuki et al., 2004; Yamagishi et al., 2003) and wingless-related (Takeuchi et al., 2003) superfamilies.

Haploinsufficiencies for several human T-box genes have been linked to congenital anomaly syndromes (Bongers et al., 2004; Packham and Brook, 2003). Two of these involve cardiac malformations. Di George syndrome, also occurring as part of chromosome 22q11 deletion syndrome, is characterized by dysmorphogenesis of the cardiac outflow tract (OFT), as well as thymic, splenic and craniofacial abnormalities (Yamagishi and Srivastava, 2003). Holt Oram

syndrome is characterized by congenital abnormalities of the upper limbs and heart, the latter involving atrial and ventricular septal defects, tetralogy of Fallot and atrioventricular conduction block (Gruber and Epstein, 2004). Targeted mutation of causative genes in mice has reproduced many aspects of the human cardiac disease phenotypes, thus providing valuable models for understanding underlying mechanisms (Bruneau et al., 2001; Lindsay et al., 2001; Merscher et al., 2001; Yamagishi and Srivastava, 2003).

*Tbx20* is an ancient member of the T-box gene subfamily related to *Tbx1* (Plageman and Yutzey, 2005). The *Drosophila* gene is expressed in early cardioblasts of the dorsal vessel of the fly, a primitive heart-like organ (Griffin et al., 2000). During fish development, *Tbx20* (*hrT*) is expressed in cardiac progenitors, developing heart and endothelial cells of the dorsal aorta (Ahn et al., 2000; Griffin et al., 2000), and *Tbx20* orthologs with analogous expression patterns have since been identified from frog, chicken, mice and humans (Carson et al., 2000; Horb and Thomsen, 1999; Kraus et al., 2001; Meins et al., 2000; Stennard et al., 2003; Yamagishi et al., 2004). During mouse development, *Tbx20* is expressed in the cardiac crescent prior to heart tube formation, then in myocardium and endocardium of the looping heart (Kraus et al., 2001; Stennard et al., 2003).

As for other T-box factors (Bruneau et al., 2001; Casey et al., 1999; Habets et al., 2002; Harrelson et al., 2004; He et al., 1999; Hoogaars et al., 2004; Hsueh et al., 2000; Kispert et al., 1995; Lamolet et al., 2001; Paxton et al., 2002; Stennard et al., 2003; Tada and Smith, 2000), *Tbx20* can regulate transcription of target genes positively and negatively, depending on the particular isoform expressed and, potentially, cellular context (Plageman and Yutzey, 2004; Stennard et al., 2003). The T-box DNA-binding domain of *Tbx20* can associate specifically, albeit weakly, with the consensus DNA half site sequence defined for the brachyury (T) protein, the founding member of the T-box family, and can interact in solution and function synergistically with homeodomain factor *Nkx2-5* and zinc finger factor *Gata4* (Stennard et al., 2003).

Morpholino oligonucleotide knockdown of *Tbx20* in fish produces small and dysmorphic hearts showing upregulation of *Tbx5*, ectopic expression of blood markers caudally and an abnormally patterned aorta (Szeto et al., 2002). *Tbx20* downregulation in frogs leads to absent (Horb and Thomsen, 1999) or dysmorphic (Brown et al., 2005) hearts. Frog *Tbx20* physically interacts with *Tbx5* and cardiac defects in embryos are more frequent and severe if both proteins are concomitantly inhibited (Brown et al., 2005). Enforced expression experiments in frog embryos show that mouse *Tbx20* can induce mesodermal and endodermal cell fates and their coordinated cell migration (Stennard et al., 2003).

We report here the loss-of-function phenotype of murine *Tbx20*. *Tbx20* null embryos showed grossly abnormal cardiac development and arrested yolk sac vascular remodeling. Our analysis has highlighted a role for *Tbx20* as a transcriptional repressor during the primary lineage split in myocardium into chamber and non-chamber fates. Furthermore, hierarchical interaction between different T-box genes was revealed as a central element of early heart patterning and morphogenesis. *Tbx20* also acts in adult heart function and homeostasis, with implications for human cardiomyopathies.

## Materials and methods

### Expression analysis

In-situ hybridization and histochemical methods were as described (Biben and Harvey, 1997; Wang et al., 2000). Rat anti-Pecam (DAKO) and rabbit anti-phosphohistone H3 (Upstate) antibodies were used at 1:200, and secondary antibodies at 1:250, with the Elite ABC kit (Vector Laboratories) and DAB substrate (Sigma). Apoptotic cells were detected using DeadEnd™ Fluorometric TUNEL System (Promega). Total RNA was isolated from hearts using Trizol (Invitrogen) and subjected to Northern analysis as described (Fatkin et al., 2000). RT-PCR using 1 µg of total RNA treated with RQ1 DNase (Promega) and 35 cycles of PCR (Rotor-Gene3000, Corbett Technologies) was as described (Stennard et al., 2003).

### Embryo culture

Explants from C57BL/6J E8.5 embryos were cultured in DMEM media containing 0.5% (v/v) heat-inactivated fetal bovine serum, 10 mmol/l glutamine, 100 units/ml penicillin and streptomycin (GibcoBRL) in 1% agarose-coated wells for 24 hours with or without recombinant human Heregulin β2 (10<sup>-9</sup> mol/l) (Fiddes et al., 1995).

### Gene targeted mice

*Tbx20<sup>lacZ/+</sup>* mice were generated by Ozgene Pty Ltd (Perth, Australia). *Nkx2-5<sup>lacZ</sup>* mice were generated using a vector similar to one described (Biben et al., 2000), in which a *lacZ* cassette was inserted in frame into exon 1 (M.S., C.B. and R.P.H., unpublished).

### Transthoracic echocardiography


Two-dimensional echocardiographic images were obtained using a Sonos 5500 ultrasonograph with 12 MHz probe (Philips Medical Systems) as described (Fatkin et al., 2000). Statistical significance of data was determined by ANOVA and Student's *t*-test.

## Results

### Generation of *Tbx20* mutant mice

To target the *Tbx20* gene by homologous recombination, we engineered a conditional vector (Fig. 1A) in which a Cre recombinase recognition site (loxP) was inserted just 5' of the methionine initiation codon in exon I, and a cassette consisting of a loxP-flanked neomycin resistance gene (*pgkneo*) followed by a nuclear-localizing β-galactosidase gene (*lacZ*) was incorporated into intron 3 (Fig. 1A). Founders were crossed with transgenic mice expressing Cre recombinase in the germline (Schwenk et al., 1995). Cre deletion of floxed *Tbx20* sequences led to excision of genomic regions encoding the methionine initiation codon, first 181 amino acids of the *Tbx20* protein and the *pgkneo* cassette (Fig. 1B-E). Deletion also brought the *lacZ* reporter directly adjacent to *Tbx20* cis-regulatory elements.

In *Tbx20<sup>lacZ/+</sup>* embryos at embryonic day (E) 7.5, *lacZ* expression was robust in the cardiac crescent, as well as amniotic and yolk sac mesoderm (Fig. 1F), confirming published in-situ hybridization patterns (Carson et al., 2000; Kraus et al., 2001; Stennard et al., 2003). Patterns at E7.5-10.5 in heart, yolk sac, brain, spinal chord, eye and allantois also confirmed published studies and revealed additional sites of expression in pharyngeal endoderm, endothelium of the dorsal aorta and associated sympathetic ganglia, vitelline and placental vessels, and adrenal medulla (Fig. 1F-I, Fig. 2E-G and data not shown).



**Fig. 1.** Generation of *Tbx20* null mice. (A) Schematic representation of the wild-type *Tbx20* locus (exons I-IV indicated as black boxes) above the *Tbx20* targeting construct and targeted *Tbx20<sup>Flox</sup>* and *Tbx20<sup>lacZ</sup>* alleles. Arrows represent the direction of transcription of neomycin (neo) and *lacZ* genes. Neo is driven by the phosphoglycerokinase promoter. Arrowheads represent loxP sites. P1 and P2 indicate 5' and 3' Southern screening probes. (B,C) Validation of the primary targeting event in embryonic stem cells using Southern analysis of *ScaI*- and *NcoI*-digested DNA and probes P1 and P2. (D) PCR genotyping assay detecting wild-type (174 bp), Floxed (220 bp) and/or Cre-deleted (302 bp) *Tbx20* alleles in mice or embryos of indicated genotypes. (E) RT-PCR analysis of RNA extracted from wild-type or *Tbx20<sup>lacZ/lacZ</sup>* embryonic hearts demonstrating deletion of exons I-III and preservation of exons IV-VI (read-through from *lacZ* cassette) in the mutant. (F-I) Expression of *lacZ* in *Tbx20<sup>lacZ/+</sup>* embryos at E7.5-10.5. am, amniotic mesoderm; cp, cardiac progenitors; da, dorsal aorta; e, eye primordium; ecc, endocardial cushion; en, endothelium; h, heart; hb, hindbrain; ias, inter-atrial septum; la, left atrium; N, *NcoI*; neo, neomycin; nt, neural tube; pe, pharyngeal endoderm; ra, right atrium; S, *ScaI*; v, ventricle; wt, wild type; ysm, yolk sac mesoderm.


### Abnormal cardiac morphogenesis in *Tbx20<sup>lacZ/lacZ</sup>* embryos

Homozygous *Tbx20<sup>lacZ/lacZ</sup>* embryos appeared normal until E7.5 but subsequently showed severe cardiac dysmorphogenesis and arrested yolk sac vascular remodeling, and all died around E10.5. Heart tube formation was retarded and abnormal from the outset. The primary ventricular chamber was small, looping was blocked and, notably, there was a significant delay in closure of the foregut pocket at caudal levels. From E8.0-9.5, a distinct additional compartment in the outflow region became progressively obvious in mutant hearts, resulting by E9.5 in a heart tube with two small chamber-like swellings separated by a circumferential sulcus (Fig. 2A-D). Apart from occasional inward myocardial protrusions, these chambers did not form trabeculae and no endocardial cushions were evident in the outflow or atrioventricular regions (Fig. 2E-J and data not shown). *LacZ* expression was largely as expected in mutant

hearts, although there was ectopic activation in dorsal mesocardial tissue in the sinuatrial region (Fig. 2G,J).

### Cardiac gene expression in *Tbx20* null embryos

The early transcription factor program was significantly compromised in *Tbx20<sup>lacZ/lacZ</sup>* hearts. Expression of T-box factor gene *Tbx5* was reduced at E8.5, although the caudal-high, graded pattern seen in normal hearts was preserved (Bruneau et al., 1999) (Fig. 3A). Expression had recovered somewhat by E9.5 (Fig. 3B), suggesting delayed activation. The cranial limit of *Tbx5* expression at E9.5 is normally at the level of the interventricular sulcus (Bruneau et al., 1999). In mutants, it was at the sulcus between the inflow and outflow chamber-like swellings (Fig. 3B), suggesting that these swellings represent precursors of the normal systemic (left) ventricle and pulmonary (right) ventricle/OFT, respectively. Consistent with this model, *Hey1*, encoding a basic helix-loop-helix factor acting downstream of Notch signaling and




**Fig. 2.** Abnormal heart morphogenesis in *Tbx20<sup>lacZ/lacZ</sup>* embryos. (A-D) Scanning electron micrographs of wild-type and *Tbx20<sup>lacZ/lacZ</sup>* embryos at E9.25. Relative scale bars are indicated. (E-J) Sections of E9.0 *Tbx20<sup>lacZ/+</sup>* and *Tbx20<sup>lacZ/lacZ</sup>* embryos stained for lacZ. Avc, atrioventricular canal; fg, foregut; iv, inflow ventricle-like chamber; la, left atrium; lv, left ventricle; oft, outflow tract; ov, outflow ventricle-like chamber; rv, ventricle; sa, sinatrium; wt, wild type.

expressed predominantly in endocardium of the OFT in normal E9.5 hearts (Iso et al., 2003), was expressed only in the outflow ventricle-like chamber in mutants (Fig. 3C).

Expression levels of the genes for homeodomain factor *Nkx2-5*, zinc finger factor *Gata4* and MADS domain factor *Mef2c* were also significantly reduced in *Tbx20<sup>lacZ/lacZ</sup>* hearts at both E8.5 and 9.5 (Fig. 3D-F and data not shown). At E8.0, *Gata4* expression was undetectable in the cranial portion of the heart progenitor field corresponding to the primary myogenic lineage, suggesting delayed activation. However, *Gata4* was already activated at this time in medial and caudal portions of the heart progenitor region (Fig. 3E), potentially occupied by a distinct cardiac progenitor population termed the secondary heart field (SHF) (Cai et al., 2003; Meilhac et al., 2004). *Tbx20* mRNA and *Tbx20-lacZ* were not substantially expressed in SHF cells dorsal to the heart once the heart tube had formed (Stennard et al., 2003) (Fig. 2E-G). SHF cells are deployed to the heart after formation of the primary heart tube and form the pulmonary ventricle and OFT, with significant contributions also to the atria. Other markers of the SHF, including *Isl1* and *Fgf8* (Cai et al., 2003), were expressed approximately normally in mutants at E8.5. *Foxh1*, encoding a potential upstream regulator of *Mef2c* in SHF (data not shown) derivatives (von Both et al., 2004) was also expressed normally (Fig. 3G). However, expression of *Tnc*, encoding the matrix protein tenascin C, while normal in mutant SHF cells dorsal and caudal to the forming heart at E8.5, failed to be maintained in the mutant outflow ventricle-like chamber (Fig. 3H).

As noted above, the outflow chamber in mutant hearts probably corresponds to precursors of the pulmonary ventricle and outflow tract of normal embryos, which are derived from the anterior SHF. To assess this further, we constructed transgenic mice bearing the human placental alkaline phosphatase gene (hPLAP) driven by an enhancer of the *Mef2c* gene, previously shown to accurately mark cells of the anterior SHF and their pulmonary ventricle and outflow derivatives (Dodou et al., 2004) (Fig. 3I). *Tbx20<sup>lacZ/lacZ</sup>* embryos carrying the transgene showed strong hPLAP staining in the outflow chamber extending



**Fig. 3.** Altered cardiac transcriptional program in *Tbx20<sup>lacZ/lacZ</sup>* embryos. (A-H) Whole-mount in-situ hybridization analysis of E8.25-9.5 wild-type or *Tbx20<sup>lacZ/lacZ</sup>* embryos showing expression of genes indicated. Arrows in B indicate cranial limit of *Tbx5* expression. Arrows in E indicate cells of the primary heart field. (I) *Tbx20<sup>lacZ/+</sup>* and *Tbx20<sup>lacZ/lacZ</sup>* embryos bearing the *Mef2c*-AHF-hPLAP transgene, stained for alkaline phosphatase activity. oft, outflow tract; wt, wild type.


caudally to the sulcus at E9.0, strongly supporting our hypothesis that this chamber is SHF-derived. Weaker staining was also seen in the caudal ventricle-like chamber, suggesting a contribution to this chamber from the anterior SHF. A minority of anterior SHF cells do contribute to the left ventricle in wild-type embryos (Cai et al., 2003), although the *Mef2c* enhancer used in these experiments is normally downregulated in those cells (D.J.M. and B.L.B., unpublished).

Excessive cell death was not detected in mutant myocardium or endocardium (data not shown). We therefore measured the mitotic index (proportion of cells expressing phosphohistone H3) in three zones of E9.0 hearts corresponding in wild-type embryos to the sinatrium, systemic ventricle and pulmonary ventricle/OFT. The myocardium of the outflow ventricle-like chamber in mutants ( $n=2$ ) had a mitotic index 6 to 7-fold less than the equivalent region in controls ( $n=2$ ;  $P<0.0001$ , chi-squared test) (see Table S1 in the supplementary material). The index in the inflow ventricle-like chamber was also reduced, although less so (2.2 and 3.3-fold;  $P=0.056$  and  $0.008$ ), while indices in the sinatrium and head mesoderm were normal.

### Expanded cardiac pre-pattern in *Tbx20* and *Tbx20/Nkx2-5* homozygotes

The *Myl2* gene, which encodes myosin light chain 2v, is expressed in ventricles and the atrioventricular canal (AVC), but not atria, betraying a molecular pre-pattern in the forming heart (Fig. 4A). How this pre-pattern is established is unknown, although maximal *Myl2* expression requires the homeodomain factor *Nkx2-5* (Lyons et al., 1995). *Myl2* was expressed in *Tbx20<sup>lacZ/lacZ</sup>* hearts at a level diminished compared with wild-type controls at E9.25, but nonetheless considerably higher than seen in *Nkx2-5<sup>GFP/GFP</sup>* embryos, which lack *Nkx2-5* function (Biben et al., 2000) (Fig. 4A). Expression encroached partially into the outflow ventricle-like chamber of *Tbx20* mutants (Fig. 4B), consistent with the notion discussed above that this chamber is SHF-derived and composed of progenitors that would normally form the pulmonary ventricle and OFT.


However, while the caudal boundary of *Myl2* expression was within the AVC in normal embryos, it was inappropriately



**Fig. 4.** Disrupted cardiac pre-pattern and *Tbx2* expression in *Tbx20<sup>lacZ/lacZ</sup>* embryos. (A–M) Whole-mount in-situ hybridization analysis of wild-type (wt), *Tbx20<sup>lacZ/lacZ</sup>*, *Nkx2-5<sup>GFP/GFP</sup>* or compound *Tbx20<sup>lacZ/lacZ</sup>/*Nkx2-5<sup>GFP/GFP</sup>** embryos using indicated probes. Panel C shows sections of whole-mount embryos at the level of the sinatrium. sa, sinatrium.

positioned within the sinatrial region of *Tbx20<sup>lacZ/lacZ</sup>* hearts (Fig. 4B), clearly evident in sections (Fig. 4C). This was even more pronounced in doubly homozygous *Tbx20<sup>lacZ/lacZ</sup>/*Nkx2-5<sup>GFP/GFP</sup>** embryos, in which *Myl2* expression extended throughout the sinus venosus (Fig. 4B). The hearts of these embryos appeared to show a combination of the abnormalities seen in single homozygotes. However, the fact that the doubly homozygous embryos ‘reactivate’ *Myl2* to levels well above those seen in single *Nkx2-5<sup>GFP/GFP</sup>* homozygotes (Fig. 4A) strongly implicates transcriptional repression at or downstream of *Tbx20* in the genetic circuitry controlling *Myl2* (see Discussion).

To investigate the apparent expansion of the *Myl2* expression domain further, we compared expression of *Myl2* and its relative *Myl7* (encoding myosin light chain 2a) at earlier stages. Despite the obvious delay in heart tube formation in mutants, dimensions of the cardiac myogenic field, as highlighted by *Myl7*, appeared comparable to those of controls at E8.5 (Fig. 4D,E). *Myl2* expression was highly regional within the *Myl7* domain. However, expression in mutants, unlike that in wild types, extended into the apparent sinatrial region. Furthermore, before



**Fig. 5.** *Tbx20* expression is repressed by Nrg1. (A–D) Panels show whole-mount in-situ hybridizations using probes indicated on E9.5 control embryos compared with embryo explants cultured from E8.5–9.5 in control medium or medium plus 10<sup>-9</sup> mol/l Nrg1.

and during the onset of heart tube formation at E7.75 and 8.0, respectively, *Myl2* expression was qualitatively different and expanded caudally and medio-anteriorly in mutants relative to somite-matched controls (Fig. 4F and data not shown). At E7.75, approximately three times the number of cells expressed *Myl2* in the mutants, demonstrating primary dysregulation of patterning processes (see Discussion).

Spatial specification of the atrial domain appeared nonetheless normal in mutants. Although diminished, a morphological sinuatrium and AVC had formed by E9.5 (Fig. 2C,D). Furthermore, the regional expression of *Hey1* was spatially correct in the outflow domain (see above) and sinuatrial region, albeit downregulated significantly in this latter domain (Fig. 4G). Expression of *Aldh1a2*, which overlaps the sinuatrial region of normal hearts and is essential for atrial specification (Niederreither et al., 2001), was also normal in *Tbx20<sup>lacZ/lacZ</sup>* embryos (Fig. 4H).

### Chamber formation in *Tbx20* mutants

The myogenic layer of the early heart tube undergoes an initial regional specialization to form working myocardium of the ventricles and atrial appendages (Christoffels et al., 2000), an event that depends on transcription factors Nkx2-5 (Palmer et al., 2001), *Tbx5* (Bruneau et al., 2001) and *Foxh1* (von Both et al., 2004). Expression levels of *Nppa* and *Smpx*, markers of chamber myocardium (Christoffels et al., 2000; Palmer et al., 2001), were severely reduced in mutant hearts at E8.5 and 9.5 (Fig. 4I–J), demonstrating lack of chamber differentiation. *Hand1*, expressed predominantly in the forming left ventricle, was also dramatically downregulated (Fig. 4K).


In the looping heart, non-chamber myocardium retains the slow conduction features evident in the primary heart tube and is destined to form elements of the central conduction system (Christoffels et al., 2004a). *Tbx2*, encoding another member of the T-box family, is expressed in non-chamber myocardium, most prominently in the AVC (Habets et al., 2002), where it has been proposed to repress formation of chamber myocardium (Christoffels et al., 2004b; Harrelson et al., 2004). By contrast to the highly regional expression of *Tbx2* in the forming AVC in normal embryos at E8.5, *Tbx2* was strikingly upregulated and ectopically expressed throughout the entire *Tbx20<sup>lacZ/lacZ</sup>* mutant heart (Fig. 4L). The pattern extended considerably more caudally in lateral plate mesoderm than in wild-type embryos, identical to the patterns of *Myl7* and *Tbx20-lacZ* at this stage (Fig. 4E and data not shown). We examined *Tbx2* expression in *Nkx2-5<sup>GFP/GFP</sup>* embryos, in which formation of chamber myocardium is also blocked (Palmer et al., 2001). *Tbx2* was expressed in the normal pattern in these embryos, although slightly diminished in level (Fig. 4L), suggesting that upregulation of *Tbx2* in all or most myogenic progenitor cells in *Tbx20<sup>lacZ/lacZ</sup>* embryos is not a default state arising from loss of chamber myocardium (see Discussion), and that *Tbx20*, directly or indirectly, represses *Tbx2* and plays a major role in its regional expression. As expected, *Tbx20<sup>lacZ/lacZ</sup>/Nkx2-5<sup>GFP/GFP</sup>* doubly homozygous embryos also showed upregulation of *Tbx2* across the heart (Fig. 4L).

*Bmp2* is expressed in myocardium of the AVC and OFT in the looping heart and has been proposed to positively regulate *Tbx2* and establish its regional pattern (Yamada et al., 2000). We therefore assessed expression of *Bmp2* and *Bmp4* in the early looping hearts of E8.0 embryos. *Bmp2* expression was in fact severely downregulated in *Tbx20<sup>lacZ/lacZ</sup>* hearts at this stage (Fig. 4M), demonstrating that *Tbx2* upregulation in *Tbx20<sup>lacZ/lacZ</sup>* hearts occurs independently of *Bmp2*. The pattern of *Bmp4* expression was normal (data not shown).

### *Tbx20* is repressed by neuregulin 1

The data above show that *Tbx20* is required for chamber differentiation, although it is unclear whether this is direct or indirect. For example, loss of chamber myocardium could result from the depressed Nkx2-5 expression (Palmer et al., 2001) or, importantly, ectopic activation of *Tbx2*, a repressor of chamber-specific gene expression (Christoffels et al., 2004b). Paradoxically, *Tbx20* may itself be a chamber repressor – the long *Tbx20a* isoform, which carries strong transcriptional activation and repression domains in its C-terminal region (Stennard et al., 2003), acts as a repressor of the chamber-specific gene *Nppa1* in vitro (Plageman and Yutzey, 2004). In embryos, *Tbx20* expression is at first enhanced in forming chamber myocardium at the outer curvature, then downregulated from E9.0, initially in the more differentiated cells of trabeculae (Stennard et al., 2003), suggesting that it is non-essential for the later stages of chamber differentiation. To clarify this issue, we asked whether *Tbx20* expression increased or decreased after treatment of myocardium in situ with a pro-chamber stimulus. Neuregulin 1 (Nrg1), a member of the epidermal growth factor family of signaling ligands, is expressed in the endocardium of the early heart tube (Garratt et al., 2003) and, along with its co-receptors ErbB2 and 4, expressed in myocardium, is essential for formation of trabeculae, a morphological feature of chamber myocardium. Excess Nrg1 induces trabecular overgrowth in vivo

**Fig. 6.** Arrested development of yolk sac vasculature in *Tbx20* null embryos. (A,B) LacZ expression in yolk sac flat mounts showing lack of mature vessels in *Tbx20<sup>lacZ/lacZ</sup>* embryos, as seen in wild type. (C,D) Sections of E9.5 *Tbx20<sup>lacZ/+</sup>* or *Tbx20<sup>lacZ/lacZ</sup>* yolk sacs showing expression of lacZ in the mesodermal layer, but not in blood cells. (E,F) *Tbx20* is expressed in yolk sac mesoderm and vessel derivatives of *Tbx20<sup>lacZ/+</sup>* embryos from E11.5-14.5. Note that expression was never observed in the endodermal layer or hematopoietic cells. (G,H) Whole-mount immunohistochemical detection of *Pecam1*, a marker of mature endothelial cells at E9.5. (I,J) TUNEL showing increased apoptosis in the mesodermal layer of E9.5 mutant yolk sacs compared with wild type. (K,L) Transmission electron microscopy of E9.5 yolk sacs showing an example of perforation of the endothelial layer (red arrowheads) in the mutant. (M) RT-PCR analysis of E9.0 yolk sacs ( $n=3$ ) for markers of hemangioblast specification (*Tal1*), angioblast specification (*Kdr*) and remodeling (*Angpt1*, *Tek*) and vessel maturation (*Acta2*, *Gja5*). Scale bar: 5  $\mu$ m. e, endoderm; ec, endothelial; wt, wild type.



and enhances myofibrillogenesis in vitro (Hertig et al., 1999). We explanted the cardiac region of wild-type E8.5 embryos and cultured them with and without Nrg1 (1 nmol/l) in low serum (0.5%) medium for 24 hours. Overall, cardiac-specific gene expression was reduced in cultured explants, for some genes dramatically (Fig. 5 and data not shown), a possible consequence of cardiac unloading. However, expression of the chamber markers *Nppa* and *Cited1* in explants was restored to approximately normal levels and the correct pattern by Nrg1. The pan-myocardial marker *Actc1* (encoding  $\alpha$ -cardiac actin) was also slightly increased (Fig. 5A,B,D). Notably, however, *Tbx20* expression was significantly repressed by Nrg1 in a dose-dependent manner, and remaining expression was mostly in endocardium (Fig. 5C and data not shown). These findings support the expression data suggesting a non-essential role for *Tbx20* in the later stages of chamber differentiation. The simplest interpretation is that chamber loss in *Tbx20<sup>lacZ/lacZ</sup>* hearts is indirect, although an early direct role for *Tbx20* in setting up the chamber program cannot be excluded. Additionally, Nrg1 may be the agent that actively represses *Tbx20* during formation of chamber myocardium in vivo.

#### Arrested vascular development in *Tbx20* mutant yolk sacs

Yolk sac vasculature remodeling was also defective in *Tbx20<sup>lacZ/lacZ</sup>* embryos. An initial vascular plexus formed, but remodeling into a mature vascular bed did not occur, as highlighted by staining for *lacZ* and vascular markers (Fig. 6A-D,G,H). *Tbx20-lacZ* was expressed only in the mesodermal layer of the yolk sac, then later in all vascular derivatives (Fig. 1F; Fig.

6C,E,F). Elevated apoptosis was detected by TUNEL assay specifically in the mesodermal layer at E9.5 and ultrastructural studies showed occasional perforation of the endothelial cell layer (Fig. 6I-L). Immunohistochemistry revealed that the endothelium in *Tbx20<sup>lacZ/lacZ</sup>* yolk sacs expressed normal levels of *Kdr* and *Pecam1*, early and late markers of differentiation, respectively (Fig. 6G,H and data not shown). Expression of several other genes involved in vascular development and remodeling were also unchanged in mutants by RT-PCR (Fig. 6M). However, *Gja5*, the gene encoding connexin 40, a gap junction protein expressed specifically in smooth muscle derivatives of yolk sac mesoderm, was strongly downregulated in mutant yolk sacs (Fig. 6M). *Gja5* is known to be regulated by T-box factors (Bruneau et al., 2001; Christoffels et al., 2004b; Stennard et al., 2003) and therefore may be directly and positively regulated by *Tbx20* during yolk sac development.

#### *Tbx20* and *Nkx2-5* genetically interact in vivo

*Tbx20* protein associates directly with the homeodomain factor *Nkx2-5* and zinc finger factors *Gata4* and *Gata5*, and these factors can function synergistically to regulate promoters of cardiac genes in vitro (Stennard et al., 2003). To test for a genetic interaction between *Tbx20* and *Nkx2-5* that would support the idea of their function in common pathways, we inter-crossed *Tbx20<sup>lacZ/+</sup>* and *Nkx2-5<sup>GFP/+</sup>* mice. A proportion of *Tbx20<sup>lacZ/+</sup>/Nkx2-5<sup>GFP/+</sup>* compound heterozygotes survived to adulthood and were apparently healthy and fertile, although substantially lower numbers than the expected 25% were found at weaning (10%;  $n=18/184$ ;  $P=0.0001$ ), suggesting partial embryonic or perinatal lethality due to structural or functional

**Table 1. Atrial septal dysmorphogenesis in adult mice**

Genotype	<i>n</i>	Normal (%)	PFO (%)	ASA (%)	PFO + ASA (%)	ASD (%)
Wild type	21	18 (85.7)	3 (14.3)	0 (0)	0 (0)	0 (0)
<i>Tbx20<sup>lacZ/+</sup></i>	20	9 (45)	6 (30)	1 (5)	4 (20)	0 (0)
<i>Nkx2-5<sup>GFP/+</sup></i>	24	2 (8.3)	12 (50)	1 (4.2)	9 (37.5)	0 (0)
<i>Tbx20<sup>lacZ/+</sup>/Nkx2-5<sup>GFP/+</sup></i>	24	3 (12.5)	4 (16.6)	1 (4.2)	12 (50)*	4 (16.7) <sup>†</sup>

PFO, patent foramen ovale; ASA, atrial septal aneurysm; ASD, atrial septal defect.

\*Increase compared with PFO+ASA in *Nkx2-5<sup>GFP/+</sup>* not significant.

<sup>†</sup>*P*<0.05 compared with all other genotypes.

malformations. Single heterozygotes were represented normally.


Humans with *NKX2.5* mutations show secundum atrial septal defect (ASD) at high penetrance (Schott et al., 1998). *Nkx2-5* heterozygous mutant mice also show ASD but only rarely (1% on C57BL/6 background), although they do manifest a spectrum of less severe atrial septal abnormalities including shortened septum primum, patent foramen ovale and atrial septal aneurysm (Biben et al., 2000). They may therefore be sensitized to mutation or downregulation of other genes involved in atrial septation. *Tbx20* may be one such gene, since it is expressed in the inter-atrial septum primum (Fig. 1G). Indeed, anatomical dissection revealed frank ASD in 16% (*n*=4/24) of *Tbx20<sup>lacZ/+</sup>/Nkx2-5<sup>GFP/+</sup>* mice, while none were found in *Tbx20<sup>lacZ/+</sup>* and *Nkx2-5<sup>GFP/+</sup>* mice (*P*<0.05) (Table 1).

### In-vivo cardiac function in *Tbx20* and *Nkx2-5* heterozygous mice

Both *Tbx20* and *Nkx2-5* were expressed in adult atrial and ventricular myocardium and interventricular septum, as judged by *lacZ* staining of hearts from *Tbx20<sup>lacZ/+</sup>* and *Nkx2-5<sup>lacZ/+</sup>* knock-in mice (Fig. 7A; see Materials and methods). We therefore examined cardiac chamber function and morphology in *Tbx20* and *Nkx2-5* single and compound heterozygous mutant embryos. Echocardiographic analysis was performed on a cohort of 28 aged-matched male mice of wild type, and single or compound heterozygous genotypes at 3-4 months of age (Table 2). Body weight and heart rate did not differ between genotypes. However, the left ventricular (LV) diastolic dimension was mildly, although significantly, increased in the *Tbx20<sup>lacZ/+</sup>* and *Tbx20<sup>lacZ/+</sup>/Nkx2-5<sup>GFP/+</sup>* genotypes (*P*=0.027; ANOVA), while wall thickness was decreased by 35 and 40%, respectively (*P*<0.001). Cardiac contractile function was also compromised, as seen by the increase in LV systolic dimension by 42 and 47% (*P*<0.001), and decrease in fractional shortening by 21 and 24% (*P*<0.001). These findings are indicative of the onset of dilated cardiomyopathy (DCM). LV parameters in *Nkx2-5<sup>GFP/+</sup>* mice were also abnormal, although less so. It is noteworthy that for all LV parameters measured there was no apparent interaction between *Tbx20* and *Nkx2-5* alleles. By contrast, an increase in left atrial dimensions, evident in all mutant genotypes, was significantly more severe in *Tbx20<sup>lacZ/+</sup>/Nkx2-5<sup>GFP/+</sup>* mice (*P*<0.001).

Based on the above findings, we examined gross cardiac morphology and gene expression in a subset of mice of each genotype selected randomly from the echocardiography cohort. After arrest in diastole, overall cardiac morphology and degree of trabeculation appeared within the normal range in single heterozygotes, despite their functional deficit. However, in three of four *Tbx20<sup>lacZ/+</sup>/Nkx2-5<sup>GFP/+</sup>* mice examined, the

right ventricle was misshapen and/or enlarged without increased wall thickness, and this appeared unrelated to the



**Fig. 7.** Dilated cardiomyopathy in adult *Tbx20<sup>lacZ/+</sup>* and *Tbx20<sup>lacZ/+</sup>/Nkx2-5<sup>lacZ/+</sup>* mice. (A) Expression of LacZ in adult *Tbx20<sup>lacZ/+</sup>* and *Nkx2-5<sup>lacZ/+</sup>* hearts. (B) Dilated right ventricle in *Tbx20<sup>lacZ/+</sup>/Nkx2-5<sup>lacZ/+</sup>* heart. (C-F) Transverse sections of a wild-type heart and one *Tbx20<sup>lacZ/+</sup>/Nkx2-5<sup>lacZ/+</sup>* heart that showed right ventricular dilation (haematoxylin and eosin staining). Arrows indicate fibrosis. (G) Northern analysis of stress and hypertrophy markers in three mice of each genotype from the echocardiographic cohort. LV, left ventricle; RV, right ventricle; wt, wild type.



**Table 2. In vivo structural and functional analysis of *Tbx20*<sup>lacZ/+</sup> and *Nkx2-5*<sup>GFP/+</sup> single and compound heterozygous adult hearts**

Parameter	Wild type (n=8)	<i>Tbx20</i> <sup>lacZ/+</sup> (n=7)	<i>Nkx2-5</i> <sup>GFP/+</sup> (n=6)	<i>Tbx20</i> <sup>lacZ/+</sup> / <i>Nkx2-5</i> <sup>GFP/+</sup> (n=7)	P value*
BW <sup>†</sup> (g)	28.9±2.4	28.0±1.4	28.1±3.0	27.3±1.4	NS
HW (g)	0.16±0.03	0.15±0.02	0.15±0.03	0.16±0.03	NS
HW/BW	5.39±0.93	5.52±0.81	5.29±0.27	6.07±0.93	NS
LVWT (mm)	0.63±0.08	0.41±0.07 <sup>‡</sup>	0.55±0.07 <sup>‡,§</sup>	0.38±0.06 <sup>‡,***</sup>	<0.001
LVDD (mm)	3.43±0.08	3.68±0.15 <sup>‡</sup>	3.57±0.26	3.67±0.18 <sup>‡</sup>	0.027
LVSD (mm)	1.35±0.05	1.92±0.31 <sup>‡</sup>	1.71±0.38 <sup>‡</sup>	1.98±0.36 <sup>‡</sup>	0.001
LVFS (%)	61±1	48±6 <sup>‡</sup>	53±8 <sup>‡</sup>	46±8 <sup>‡</sup>	<0.001
LAD (mm)	1.61±0.04	1.87±0.10 <sup>‡</sup>	1.74±0.11 <sup>‡,§</sup>	2.10±0.19 <sup>‡,§,¶,***</sup>	<0.001
HR (bpm)	624±26	646±20	658±30 <sup>‡</sup>	604±79	NS

\*P value determined by ANOVA.

<sup>†</sup>Mice aged 12-18 weeks.

<sup>‡</sup>P<0.05 compared with wild type.

<sup>§</sup>P<0.05 *Tbx20*<sup>lacZ/+</sup> compared with *Nkx2-5*<sup>GFP/+</sup>.

<sup>¶</sup>P<0.05 *Tbx20*<sup>lacZ/+</sup> compared with *Tbx20*<sup>lacZ/+</sup>/*Nkx2-5*<sup>GFP/+</sup>.

<sup>\*\*\*</sup>P<0.05 *Nkx2-5*<sup>GFP/+</sup> compared with *Tbx20*<sup>lacZ/+</sup>/*Nkx2-5*<sup>GFP/+</sup>.

BW, body weight; HW, heart weight; LV, left ventricle; WT, wall thickness; DD, end-diastolic diameter; SD, end-systolic diameter; FS, fractional shortening (determined as LVDD-LVSD/LVDD); LAD, left atrial diameter; HR, heart rate; NS, not significant.

presence of ASD (Fig. 7B). Histology on transverse sections from one of the three such affected hearts showed pronounced myocyte disarray and patches of fibrosis specifically in RV myocardium (Fig. 7C-F), although examination of additional mice showed that gross myocyte disarray was not a general feature of the doubly heterozygous hearts showing RV dilation.

Despite reduced contractile function, compensatory myocardial hypertrophy was not evident in any of the mutant hearts. Specifically, there was no change in heart weights or heart weight/body weight ratios (Table 2), and no overt signs of myofiber hypertrophy. Moreover, while northern analysis showed upregulation of *Nppa*, a general marker of myocardial stress, in *Tbx20*<sup>lacZ/+</sup> hearts, multiple markers of cardiac hypertrophy were either normally expressed or diminished in mutant genotypes (Fig. 7G). *Nppa* was significantly downregulated in *Nkx2-5*<sup>GFP/+</sup> and *Tbx20*<sup>lacZ/+</sup>/*Nkx2-5*<sup>GFP/+</sup> hearts, probably because *Nkx2-5* plays a direct role in *Nppa* transcription (Durocher et al., 1996).

## Discussion

Nearly 1% of live born humans have some form of structural malformation of the heart. Recent data suggest that a few percent of these are caused by mutations in cardiac transcription factors acting in development, including T-box factors *Tbx5* and *Tbx1* (Garg et al., 2003; Prall et al., 2002; Yagi et al., 2003). In this paper, we address the function of cardiac T-box factor *Tbx20*. Our data demonstrate a central role for *Tbx20* in patterning and chamber formation in the embryonic heart via positive and negative influences on other T-box genes. We have also uncovered a role for *Tbx20* in the adult heart, where it contributes to ventricular function, integrity and adaptation.

### Morphology and gene expression in *Tbx20* mutant hearts

In the absence of *Tbx20*, heart tube development was severely retarded. By E9.5, a small, forward looped, hourglass-shaped heart with two distinct ventricle-like chambers separated by a pronounced sulcus and a diminished sinatrium had formed.

Our morphological findings are generally consistent with the consequences of *Tbx20* morpholino knockdown experiments in fish and frogs, which showed small, unlooped and dysmorphic hearts with poor chamber discrimination (Brown et al., 2005; Szeto et al., 2002).

While the cardiac progenitor field appeared to form normally in *Tbx20* mutants, there was initially a profound delay in incorporation of progenitors into the forming primary heart tube. We have previously shown that injection of *Tbx20* mRNA into *Xenopus* embryos induces mesodermal and endodermal cell fates and their coordinated migration via cell non-autonomous mechanisms (Stennard et al., 2003), and numerous other T-box factors have been implicated in control of cell migration at gastrulation (Russ et al., 2000; Tada and Smith, 2000; Yamamoto et al., 1998).

While *Tbx20* could control cell migration of cardiac progenitors directly, the effects may in part be indirect due to blocked or delayed differentiation of myocardium or endocardium. Indeed, in *Tbx20*<sup>lacZ/lacZ</sup> mutants, the early cardiac regulatory program involving transcription factors *Nkx2-5*, *Gata4* and *Mef2c* and cardiac inducing factor *Bmp2* was significantly compromised. Furthermore, expression of *Tbx20-lacZ* in foregut flags the possibility that *Tbx20* could also regulate differentiation of endoderm, a well-known source of factors that induce and support differentiation (and migration) of the cardiomyocyte lineage (Nascone and Mercola, 1996).

Development of SHF derivatives in *Tbx20*<sup>lacZ/lacZ</sup> mutants was also defective. Abnormal deployment or differentiation of SHF cells is likely to underlie conotruncal and other congenital heart defects in humans, with haploinsufficiency for the T-box factor gene, *TBX1*, expressed in SHF cells and associated endoderm, thought to be the major determinant of a spectrum of heart as well as branchial region defects associated with chromosome 22q11 deletion syndrome (Yagi et al., 2003). The outflow ventricle-like chamber in *Tbx20*<sup>lacZ/lacZ</sup> hearts is likely to be derived from the SHF, and this is supported by strong expression of the *Mef2c-SHF-hPLAP* transgene in this chamber. However, cell proliferation in the outflow chamber was severely compromised and the structure remained bulbous

and did not undergo elongation or looping as in normal hearts. Furthermore, expression of *Tnc*, encoding a matrix protein with broad regulatory functions on cell adhesion, migration and proliferation through interactions with other matrix components and cell surface receptors (Jones and Jones, 2000), was not maintained in the outflow region. Thus, *Tbx20* is essential for cell proliferation and gene expression in SHF cells once they enter the outflow region of the heart, effects that may be mediated by alterations to the extracellular matrix. This would affect clonal growth patterns in the forming heart that support correct cardiac looping and outflow tract morphogenesis (Meilhac et al., 2004). It is noteworthy that other T-box genes have been implicated in control of cell proliferation (Hatcher et al., 2001; Xu et al., 2004).

### Chamber formation and transcriptional repression in the developing heart

Chamber muscle becomes evident early in heart development from the regional expression of several genes, most notably *Nppa* and *Smpx*, and formation of trabeculae (Christoffels et al., 2000; Palmer et al., 2001). Transcription factors *Tbx5*, *Nkx2-5* and *Foxh1* are essential for its specification (Bruneau et al., 2001; Lyons et al., 1995; von Both et al., 2004), and *Tbx5* and *Nkx2-5* directly regulate chamber-specific genes in vitro (Bruneau et al., 2001; Stennard et al., 2003). In *Tbx20<sup>lacZ/lacZ</sup>* hearts, expression of chamber-specific markers was severely downregulated, indicating that they do not differentiate chamber muscle. The *Nrg1* pathway, which is necessary (but not sufficient) for chamber differentiation, repressed *Tbx20* expression in situ, suggesting that loss of chamber myocardium in *Tbx20<sup>lacZ/lacZ</sup>* hearts is indirect, probably a consequence of ectopic activation of *Tbx2*. It is still feasible, however, that *Tbx20* plays a direct positive role at the earliest stages of chamber formation.

A key finding of this work is that *Tbx2* was ectopically expressed in all or most committed myocyte progenitors in *Tbx20* mutant hearts. *Tbx2* is expressed normally in non-chamber myocardium, and in the AVC it is thought to compete with *Tbx5* for interaction with *Nkx2-5* on the *cis*-regulatory elements of chamber-specific genes, thus inhibiting their expression (Habets et al., 2002). The global expression of *Tbx2* in mutant hearts could merely reflect the loss of chamber myocardium and expansion of non-chamber myocardium. However, two facts argue against this possibility. First, *Tbx2* was markedly upregulated (3-fold by RT-PCR quantitation) as well as ectopically expressed. Second, *Tbx2* was expressed normally in the hearts of *Nkx2-5<sup>GFP/GFP</sup>* embryos, in which chamber differentiation is also blocked at the level of a controlling transcription factor. We conclude that *Tbx20* directly or indirectly represses *Tbx2* in myocardium, and that *Tbx20* plays a defining role in specification of chamber and non-chamber myocardium, a lineage digression in the early heart upon which all subsequent morphogenesis depends.

Our data suggest a model in which chamber formation in the heart involves 'default repression', a feature of virtually all well-studied, conserved, signal-induced transcriptional regulatory systems acting in development (Barolo and Posakony, 2002). Default repression occurs when a developmental process is actively repressed in the absence of its inducing signal to prevent cryptic activation by other positive factors involved in specificity. Thus, specification of

the *Tbx2* pattern in the AVC and other regions of non-chamber myocardium must involve regional and presumably signal-dependent inhibition of the repressive role of *Tbx20* on *Tbx2* expression, a possible role for Bmps (Yamada et al., 2000).

A repressive role for *Tbx20* was also evident in regulation of the *Myl2* cardiac pre-pattern. *Myl2* is expressed at only very low levels in hearts of *Nkx2-5* null embryos (Biben et al., 2000; Lyons et al., 1995), yet was 'reactivated' in *Tbx20<sup>lacZ/lacZ</sup>/Nkx2-5<sup>GFP/GFP</sup>* embryos, which lack both *Nkx2-5* and *Tbx20* function. This finding inextricably implicates transcriptional repression involving *Tbx20* in the regulation of *Myl2*. The *Myl2* pattern was also broader relative to morphological landmarks in *Tbx20<sup>lacZ/lacZ</sup>* and *Tbx20<sup>lacZ/lacZ</sup>/Nkx2-5<sup>GFP/GFP</sup>* hearts. Consistent with these findings, expanded expression of the *cmlc2* gene (an *Myl2* ortholog) into the atria was noted after morpholino knockdown of zebrafish *Tbx20*, and the ventricle-specific gene *vmhc* was also activated in this region (Szeto et al., 2002). These patterns probably reflect loss of repressive roles for *Tbx20*, and it is noteworthy that expansion of the expression domains of developmental genes, as seen here, is also one hallmark of loss of default repression (Barolo and Posakony, 2002). Our data suggest multiple repressive functions for *Tbx20* in the core cardiac regulatory program. We envisage a genetic circuitry for heart development based on interactions between multiple cardiac T-box factors and their co-factors, involving overlapping steps of repression and de-repression.

### A role for cardiac transcription factors in adult heart pathology

Our studies have also revealed a key role for *Tbx20* in adult cardiac function. *Tbx20* haploinsufficiency led to LV dilation, decreased wall thickness and contractile dysfunction, indicative of DCM. Gross dilation was also seen in the RV in some *Tbx20<sup>lacZ/+</sup>/Nkx2-5<sup>GFP/+</sup>* mice. In the atrial compartment, ASD was evident in 16% of *Tbx20<sup>lacZ/+</sup>/Nkx2-5<sup>GFP/+</sup>* mice, and there was left atrial dilation in all mutant genotypes analyzed, although most severely in the *Tbx20<sup>lacZ/+</sup>/Nkx2-5<sup>GFP/+</sup>* mice. The specific roles for *Tbx20* in adult cardiac structure and function remain to be determined. ASD is developmental in origin and our data highlight *TBX20* as a candidate ASD gene in humans. In relation to ventricular defects, we found no deficit in expression of developmental genes such as *Tbx5*, *Nkx2-5*, *Smpx* and *Gjal* in *Tbx20<sup>lacZ/+</sup>* mice (data not shown). Nevertheless, it will be important to explore the timing of onset of LV DCM to determine whether it is also developmental in origin or reflects specific adult functions for *Tbx20*. In humans, a large number of disease genes for familial DCM have been identified, including those for sarcomeric, cytoskeletal, nuclear and calcium handling proteins (Fatkin and Graham, 2002). However, known disease genes account for only a small proportion of all familial cases. Mutations in cardiac transcription factor genes may prove to be another cause of DCM in humans.

The onset of LV dilation and contractile dysfunction in *Tbx20<sup>lacZ/+</sup>* mice in the absence of hypertrophy, fingers *Tbx20* as an essential gene in the adult cardiac adaptive response. In most models of adult cardiomyopathy, hypertrophy is a component of the pathophysiological response, although it can be bypassed if structural proteins, potential sensors of biomechanical stress, are absent (Brancaccio et al., 2003; Knoll

et al., 2002). Recent work highlighting the developmental transcription factor Gata4 as a convergence point for cardiac hypertrophy pathways (Liang and Molkenkin, 2002), has supported the long-held view that developmental pathways are reactivated in hypertrophy, although mechanistic understanding is still limited and distinctions between adaptive and pathological hypertrophy are unclear (Fatkin and Graham, 2002). Further analysis of the *Tbx20* model should advance our understanding of the important link between development and adaptive responses in the adult organ.

This work was supported by a grant from the National Health and Medical Research Council (NHMRC) of Australia (256302). OWJP was a NHMRC Peter Doherty Fellow, SLD a Pfizer Foundation Australia Senior Research Fellow and DF a Sylvia and Charles Viertel Charitable Foundation Fellow. We thank Marianne Beales for help with production of transgenic mice and Eric Schmied for animal husbandry.

### Supplementary material

Supplementary material for this article is available at <http://dev.biologists.org/cgi/content/full/132/10/2451/DC1>

## References

- Ahn, D. G., Ruvinsky, I., Oates, A. C., Silver, L. M. and Ho, R. K. (2000). *tbx20*, a new vertebrate T-box gene expressed in the cranial motor neurons and developing cardiovascular structures in zebrafish. *Mech. Dev.* **95**, 253-258.
- Barolo, S. and Posakony, J. W. (2002). Three habits of highly effective signaling pathways: principles of transcriptional control by developmental cell signaling. *Genes Dev.* **16**, 1167-1181.
- Biben, C. and Harvey, R. P. (1997). Homeodomain factor Nkx2-5 controls left/right asymmetric expression of bHLH gene *eHand* during murine heart development. *Genes Dev.* **11**, 1357-1369.
- Biben, C., Weber, R., Kesteven, S., Stanley, E., McDonald, L., Elliott, D. A., Barnett, L., Koentgen, F., Robb, L., Feneley, M. et al. (2000). Cardiac septal and valvular dysmorphogenesis in mice heterozygous for mutations in the homeobox gene *Nkx2-5*. *Circ. Res.* **87**, 888-895.
- Bongers, E. M., Duijff, P. H., van Beersom, S. E., Schoots, J., van Kampen, A., Burekhardt, A., Hamel, B. C., Losan, F., Hoefsloot, L. H., Yntema, H. G. et al. (2004). Mutations in the human *TBX4* gene cause small patella syndrome. *Am. J. Hum. Genet.* **74**, 1239-1248.
- Brancaccio, M., Fratta, L., Notte, A., Hirsch, E., Poulet, R., Guazzone, S., de Acetis, M., Vecchione, C., Marino, G., Altruda, F. et al. (2003). Melusin, a muscle-specific integrin beta1-interacting protein, is required to prevent cardiac failure in response to chronic pressure overload. *Nat. Med.* **9**, 68-75.
- Brown, C. B., Wenning, J. M., Lu, M. M., Epstein, D. J., Meyers, E. N. and Epstein, J. A. (2004). Cre-mediated excision of *Fgf8* in the *Tbx1* expression domain reveals a critical role for *Fgf8* in cardiovascular development in the mouse. *Dev. Biol.* **267**, 190-202.
- Brown, D. D., Martz, S. N., Binder, O., Goetz, S. C., Price, B. M., Smith, J. C. and Conlon, F. L. (2005). *Tbx5* and *Tbx20* act synergistically to control vertebrate heart morphogenesis. *Development* **132**, 553-563.
- Bruneau, B. G., Logan, M., Davis, N., Levi, T., Tabin, C. J., Seidman, J. G. and Seidman, C. E. (1999). Chamber-specific cardiac expression of *Tbx5* and heart defects in Holt-Oram syndrome. *Dev. Biol.* **211**, 100-108.
- Bruneau, B. G., Nemer, G., Schmitt, J. P., Charron, F., Robitaille, L., Caron, S., Conner, D. A., Gessler, M., Nemer, M., Seidman, C. E. et al. (2001). A murine model of Holt-Oram syndrome defines roles of the T-box transcription factor *Tbx5* in cardiogenesis and disease. *Cell* **106**, 709-721.
- Cai, C. L., Liang, X., Shi, Y., Chu, P. H., Pfaff, S. L., Chen, J. and Evans, S. (2003). *Isl1* identifies a cardiac progenitor population that proliferates prior to differentiation and contributes a majority of cells to the heart. *Dev. Cell* **5**, 877-889.
- Carson, C. T., Kinzler, E. R. and Parr, B. A. (2000). *Tbx12*, a novel T-box gene, is expressed during early stages of heart and retinal development. *Mech. Dev.* **96**, 137-140.
- Casey, E. S., Tada, M., Fairclough, L., Wylie, C. C., Heasman, J. and Smith, J. C. (1999). *Bix4* is activated directly by VegT and mediates endoderm formation in *Xenopus* development. *Development* **126**, 4193-4200.
- Chapman, D. L. and Papaioannou, V. E. (1998). Three neural tubes in mouse embryos with mutations in the T-box gene *Tbx6*. *Nature* **391**, 695-697.
- Christoffels, V. M., Habets, P. E., Franco, D., Campione, M., de Jong, F., Lamers, W. H., Bao, Z. Z., Palmer, S., Biben, C., Harvey, R. P. et al. (2000). Chamber formation and morphogenesis in the developing mammalian heart. *Dev. Biol.* **223**, 266-278.
- Christoffels, V. M., Burch, J. B. and Moorman, A. F. (2004a). Architectural Plan for the Heart: Early Patterning and Delineation of the Chambers and the Nodes. *Trends Cardiovasc. Med.* **14**, 301-307.
- Christoffels, V. M., Hoogaars, W. M., Tessari, A., Clout, D. E., Moorman, A. F. and Campione, M. (2004b). T-box transcription factor *Tbx2* represses differentiation and formation of the cardiac chambers. *Dev. Dyn.* **229**, 763-770.
- Cripps, R. M. and Olson, E. N. (2002). Control of cardiac development by an evolutionarily conserved transcriptional network. *Dev. Biol.* **246**, 14-28.
- Dodou, E., Verzi, M. P., Anderson, J. P., Xu, S. M. and Black, B. L. (2004). *Mef2c* is a direct transcriptional target of *ISL1* and *GATA* factors in the anterior heart field during mouse embryonic development. *Development* **131**, 3931-3942.
- Durocher, D., Chen, C. Y., Ardati, A., Schwartz, R. J. and Nemer, M. (1996). The atrial natriuretic factor promoter is a downstream target for *Nkx-2.5* in the myocardium. *Mol. Cell Biol.* **16**, 4648-4655.
- Fatkin, D. and Graham, R. M. (2002). Molecular mechanisms of inherited cardiomyopathies. *Physiol. Rev.* **82**, 945-980.
- Fatkin, D., McConnell, B. K., Mudd, J. O., Semsarian, C., Moskowitz, I. G., Schoen, F. J., Giewat, M., Seidman, C. E. and Seidman, J. G. (2000). An abnormal Ca(2+) response in mutant sarcomere protein-mediated familial hypertrophic cardiomyopathy. *J. Clin. Invest.* **106**, 1351-1359.
- Fiddes, R. J., Janes, P. W., Sanderson, G. M., Sivertsen, S. P., Sutherland, R. L. and Daly, R. J. (1995). Heregulin (HRG)-induced mitogenic signaling and cytotoxic activity of a HRG/PE40 ligand toxin in human breast cancer cells. *Cell Growth Differ.* **6**, 1567-1577.
- Garg, V., Kathiriyi, I. S., Barnes, R., Schluterman, M. K., King, I. N., Butler, C. A., Rothrock, C. R., Eapen, R. S., Hirayama-Yamada, K., Joo, K. et al. (2003). *GATA4* mutations cause human congenital heart defects and reveal an interaction with *TBX5*. *Nature* **424**, 443-447.
- Garratt, A. N., Ozcelik, C. and Birchmeier, C. (2003). *ErbB2* pathways in heart and neural diseases. *Trends Cardiovasc. Med.* **13**, 80-86.
- Griffin, K. J., Stoller, J., Gibson, M., Chen, S., Yelon, D., Stainier, D. Y. and Kimelman, D. (2000). A conserved role for H15-related T-box transcription factors in zebrafish and *Drosophila* heart formation. *Dev. Biol.* **218**, 235-247.
- Gruber, P. J. and Epstein, J. A. (2004). Development gone awry: congenital heart disease. *Circ. Res.* **94**, 273-283.
- Habets, P. E., Moorman, A. F., Clout, D. E., van Roon, M. A., Lingbeek, M., van Lohuizen, M., Campione, M. and Christoffels, V. M. (2002). Cooperative action of *Tbx2* and *Nkx2.5* inhibits ANF expression in the atrioventricular canal: implications for cardiac chamber formation. *Genes Dev.* **16**, 1234-1246.
- Harrelson, Z., Kelly, R. G., Goldin, S. N., Gibson-Brown, J. J., Bollag, R. J., Silver, L. M. and Papaioannou, V. E. (2004). *Tbx2* is essential for patterning the atrioventricular canal and for morphogenesis of the outflow tract during heart development. *Development* **131**, 5041-5052.
- Harvey, R. P. (2002). Molecular Determinants of Cardiac Development and Congenital Disease. In *Mouse Development: Patterning, Morphogenesis, and Organogenesis* (ed. J. Rossant and P. P. L. Tam), pp. 331-370. San Diego: Academic Press.
- Hatcher, C. J., Kim, M. S., Mah, C. S., Goldstein, M. M., Wong, B., Mikawa, T. and Basson, C. T. (2001). *TBX5* transcription factor regulates cell proliferation during cardiogenesis. *Dev. Biol.* **230**, 177-188.
- He, M., Wen, L., Campbell, C. E., Wu, J. Y. and Rao, Y. (1999). Transcription repression by *Xenopus* ET and its human ortholog *TBX3*, a gene involved in ulnar-mammary syndrome. *Proc. Natl. Acad. Sci. USA* **96**, 10212-10217.
- Hertig, C. M., Kubalak, S. W., Wang, Y. and Chien, K. R. (1999). Synergistic roles of neuregulin-1 and insulin-like growth factor-I in activation of the phosphatidylinositol 3-kinase pathway and cardiac chamber morphogenesis. *J. Biol. Chem.* **274**, 37362-37369.
- Hoogaars, W. M., Tessari, A., Moorman, A. F., de Boer, P. A., Hagoort, J., Soufan, A. T., Campione, M. and Christoffels, V. M. (2004). The

- transcriptional repressor Tbx3 delineates the developing central conduction system of the heart. *Cardiovasc. Res.* **62**, 489-499.
- Horb, M. E. and Thomsen, G. H.** (1999). Tbx5 is essential for heart development. *Development* **126**, 1739-1751.
- Hsueh, Y. P., Wang, T. F., Yang, F. C. and Sheng, M.** (2000). Nuclear translocation and transcription regulation by the membrane-associated guanylate kinase CASK/LIN-2. *Nature* **404**, 298-302.
- Hu, T., Yamagishi, H., Maeda, J., McAnally, J., Yamagishi, C. and Srivastava, D.** (2004). Tbx1 regulates fibroblast growth factors in the anterior heart field through a reinforcing autoregulatory loop involving forkhead transcription factors. *Development* **131**, 5491-5502.
- Iso, T., Kedes, L. and Hamamori, Y.** (2003). HES and HERP families: multiple effectors of the Notch signaling pathway. *J. Cell Physiol.* **194**, 237-255.
- Jones, F. S. and Jones, P. L.** (2000). The tenascin family of ECM glycoproteins: structure, function, and regulation during embryonic development and tissue remodeling. *Dev. Dyn.* **218**, 235-259.
- Kispert, A., Koschorz, B. and Herrmann, B. G.** (1995). The T protein encoded by Brachyury is a tissue-specific transcription factor. *EMBO J.* **14**, 4763-4772.
- Knoll, R., Hoshijima, M., Hoffman, H. M., Person, V., Lorenzen-Schmidt, I., Bang, M. L., Hayashi, T., Shiga, N., Yasukawa, H., Schaper, W. et al.** (2002). The cardiac mechanical stretch sensor machinery involves a Z disc complex that is defective in a subset of human dilated cardiomyopathy. *Cell* **111**, 943-955.
- Kraus, F., Haenig, B. and Kispert, A.** (2001). Cloning and expression analysis of the mouse T-box gene tbx20. *Mech. Dev.* **100**, 87-91.
- Lamolet, B., Pulichino, A. M., Lamonerie, T., Gauthier, Y., Brue, T., Enjalbert, A. and Drouin, J.** (2001). A pituitary cell-restricted T box factor, Tpit, activates POMC transcription in cooperation with Pitx homeoproteins. *Cell* **104**, 849-859.
- Liang, Q. and Molkentin, J. D.** (2002). Divergent signaling pathways converge on GATA4 to regulate cardiac hypertrophic gene expression. *J. Mol. Cell. Cardiol.* **34**, 611-616.
- Lindsay, E. A., Vitelli, F., Su, H., Morishima, M., Huynh, T., Pramparo, T., Jurecic, V., Ogunrinu, G., Sutherland, H. F., Scambler, P. J. et al.** (2001). Tbx1 haploinsufficiency in the DiGeorge syndrome region causes aortic arch defects in mice. *Nature* **410**, 97-101.
- Lyons, I., Parsons, L. M., Hartley, L., Li, R., Andrews, J. E., Robb, L. and Harvey, R. P.** (1995). Myogenic and morphogenetic defects in the heart tubes of murine embryos lacking the homeo box gene Nkx2-5. *Genes Dev.* **9**, 1654-1666.
- Meilhac, S. M., Esner, M., Kelly, R. G., Nicolas, J. F. and Buckingham, M. E.** (2004). The clonal origin of myocardial cells in different regions of the embryonic mouse heart. *Dev. Cell* **6**, 685-698.
- Meins, M., Henderson, D. J., Bhattacharya, S. S. and Sowden, J. C.** (2000). Characterization of the human TBX20 gene, a new member of the T-Box gene family closely related to the Drosophila H15 gene. *Genomics* **67**, 317-332.
- Merscher, S., Funke, B., Epstein, J. A., Heyer, J., Puech, A., Lu, M. M., Xavier, R. J., Demay, M. B., Russell, R. G., Factor, S. et al.** (2001). TBX1 is responsible for cardiovascular defects in velo-cardio-facial/DiGeorge syndrome. *Cell* **104**, 619-629.
- Naiche, L. A. and Papaioannou, V. E.** (2003). Loss of Tbx4 blocks hindlimb development and affects vascularization and fusion of the allantois. *Development* **130**, 2681-2693.
- Nascone, N. and Mercola, M.** (1996). Endoderm and Cardiogenesis: New Insights. *Trends Cardio. Med.* **6**, 211-216.
- Niederreither, K., Vermot, J., Messaddeq, N., Schuhbaur, B., Chambon, P. and Dolle, P.** (2001). Embryonic retinoic acid synthesis is essential for heart morphogenesis in the mouse. *Development* **128**, 1019-1031.
- Packham, E. A. and Brook, J. D.** (2003). T-box genes in human disorders. *Hum. Mol. Genet.* **12**, R37-R44.
- Palmer, S., Groves, N., Schindeler, A., Yeoh, T., Biben, C., Wang, C. C., Sparrow, D. B., Barnett, L., Jenkins, N. A., Copeland, N. G. et al.** (2001). The small muscle-specific protein Csl modifies cell shape and promotes myocyte fusion in an insulin-like growth factor I-dependent manner. *J. Cell Biol.* **153**, 985-998.
- Paxton, C., Zhao, H., Chin, Y., Langner, K. and Reecy, J.** (2002). Murine Tbx2 contains domains that activate and repress gene transcription. *Gene* **283**, 117-124.
- Plageman, T. F., Jr and Yutzey, K. E.** (2004). Differential expression and function of Tbx5 and Tbx20 in cardiac development. *J. Biol. Chem.* **279**, 19026-19034.
- Plageman, T. F., Jr and Yutzey, K. E.** (2005). T-box genes and heart development: Putting the "T" in heart. *Dev. Dyn.* **232**, 11-20.
- Prall, O. W., Elliott, D. A. and Harvey, R. P.** (2002). Developmental paradigms in heart disease: insights from tinman. *Ann. Med.* **34**, 148-156.
- Russ, A. P., Wattler, S., Colledge, W. H., Aparicio, S. A., Carlton, M. B., Pearce, J. J., Barton, S. C., Surani, M. A., Ryan, K., Nehls, M. C. et al.** (2000). Eomesodermin is required for mouse trophoblast development and mesoderm formation. *Nature* **404**, 95-99.
- Sakiyama, J., Yamagishi, A. and Kuroiwa, A.** (2003). Tbx4-Fgf10 system controls lung bud formation during chicken embryonic development. *Development* **130**, 1225-1234.
- Schott, J. J., Benson, D. W., Basson, C. T., Pease, W., Silberbach, G. M., Moak, J. P., Maron, B. J., Seidman, C. E. and Seidman, J. G.** (1998). Congenital heart disease caused by mutations in the transcription factor NKX2-5. *Science* **281**, 108-111.
- Schwenk, F., Baron, U. and Rajewsky, K.** (1995). A cre-transgenic mouse strain for the ubiquitous deletion of loxP-flanked gene segments including deletion in germ cells. *Nucleic Acids Res.* **23**, 5080-5081.
- Smith, J.** (1999). T-box genes: what they do and how they do it. *Trends Genet.* **15**, 154-158.
- Stennard, F. A., Costa, M. W., Elliott, D. A., Rankin, S., Haast, S. J., Lai, D., McDonald, L. P., Niederreither, K., Dolle, P., Bruneau, B. G. et al.** (2003). Cardiac T-box factor Tbx20 directly interacts with Nkx2-5, GATA4, and GATA5 in regulation of gene expression in the developing heart. *Dev. Biol.* **262**, 206-224.
- Suzuki, T., Takeuchi, J., Koshiba-Takeuchi, K. and Ogura, T.** (2004). Tbx Genes Specify Posterior Digit Identity through Shh and BMP Signaling. *Dev. Cell* **6**, 43-53.
- Szeto, D. P., Griffin, K. J. and Kimelman, D.** (2002). HrT is required for cardiovascular development in zebrafish. *Development* **129**, 5093-5101.
- Tada, M. and Smith, J. C.** (2000). Xwnt11 is a target of Xenopus Brachyury: regulation of gastrulation movements via Dishevelled, but not through the canonical Wnt pathway. *Development* **127**, 2227-2238.
- Takeuchi, J. K., Koshiba-Takeuchi, K., Suzuki, T., Kamimura, M., Ogura, K. and Ogura, T.** (2003). Tbx5 and Tbx4 trigger limb initiation through activation of the Wnt/Fgf signaling cascade. *Development* **130**, 2729-2739.
- von Both, I., Silvestri, C., Erdemir, T., Lickert, H., Walls, J. R., Henkelman, R. M., Rossant, J., Harvey, R. P., Attisano, L. and Wrana, J. L.** (2004). Foxh1 is essential for development of the anterior heart field. *Dev. Cell* **7**, 331-345.
- Wang, C. C., Biben, C., Robb, L., Nassir, F., Barnett, L., Davidson, N. O., Koentgen, F., Tarlinton, D. and Harvey, R. P.** (2000). Homeodomain factor Nkx2-3 controls regional expression of leukocyte homing coreceptor MadCAM-1 in specialized endothelial cells of the viscera. *Dev. Biol.* **224**, 152-167.
- Xu, H., Morishima, M., Wylie, J. N., Schwartz, R. J., Bruneau, B. G., Lindsay, E. A. and Baldini, A.** (2004). Tbx1 has a dual role in the morphogenesis of the cardiac outflow tract. *Development* **131**, 3217-3227.
- Yagi, H., Furutani, Y., Hamada, H., Sasaki, T., Asakawa, S., Minoshima, S., Ichida, F., Joo, K., Kimura, M., Imamura, S. et al.** (2003). Role of TBX1 in human del22q11.2 syndrome. *Lancet* **362**, 1366-1373.
- Yamada, M., Revelli, J. P., Eichele, G., Barron, M. and Schwartz, R. J.** (2000). Expression of chick Tbx-2, Tbx-3, and Tbx-5 genes during early heart development: evidence for BMP2 induction of Tbx2. *Dev. Biol.* **228**, 95-105.
- Yamagishi, H. and Srivastava, D.** (2003). Unraveling the genetic and developmental mysteries of 22q11 deletion syndrome. *Trends Mol. Med.* **9**, 383-389.
- Yamagishi, H., Maeda, J., Hu, T., McAnally, J., Conway, S. J., Kume, T., Meyers, E. N., Yamagishi, C. and Srivastava, D.** (2003). Tbx1 is regulated by tissue-specific forkhead proteins through a common Sonic hedgehog-responsive enhancer. *Genes Dev.* **17**, 269-281.
- Yamagishi, T., Nakajima, Y., Nishimatsu, S., Nohno, T., Ando, K. and Nakamura, H.** (2004). Expression of tbx20 RNA during chick heart development. *Dev. Dyn.* **230**, 576-580.
- Yamamoto, A., Amacher, S. L., Kim, S. H., Geissert, D., Kimmel, C. B. and de Robertis, E. M.** (1998). Zebrafish paraxial protocadherin is a downstream target of spadetail involved in morphogenesis of gastrula mesoderm. *Development* **125**, 3389-3397.
- Zaffran, S. and Frasch, M.** (2002). Early signals in cardiac development. *Circ. Res.* **91**, 457-469.

**Table S1. Cell proliferation in E9.0 embryonic hearts**

Genotype (somites)	Tissue	Cells counted	Mitotic index*	<i>P</i>
Wild type (15)	SV/A	703	4.13	
	LV	731	3.56	
	RV/OFT	672	3.87	
	PM	503	6.16	
<i>Tbx20<sup>lacZ/lacZ</sup></i> (15)	SV/A	1031	2.72	NS
	IV	566	1.06	<0.01
	OV	900	0.67	< 0.0001
	PM	460	5.44	NS
Wild type (17)	SV/A	685	2.77	
	LV	568	3.35	
	RV/OFT	649	3.85	
	PM	781	4.87	
<i>Tbx20<sup>lacZ/lacZ</sup></i> (17)	SV/A	337	2.97	NS
	IV	720	1.53	=0.056
	OV	1229	0.49	<0.0001
	PM	638	4.08	NS

The analysis was performed on two wild-type and two *Tbx20<sup>lacZ/lacZ</sup>* embryos.

*P* values are with respect to a somite number-matched control using  $\chi^2$  test.

\*Percentage of DAPI stained nuclei immunopositive for phosphorylated histone H3.

A, atria; IV, inflow ventricle-like chamber; LV, left ventricle; OFT, outflow tract; OV, outflow ventricle-like chamber; PM, paraxial mesoderm; RV: right ventricle; SV, sinus venosa; NS, not significant.